

Higgs Masses in the Four Generation MSSM

Sean Litsey*

Department of Physics, Yale University, New Haven, CT 06511-8499 USA

Marc Sher†

Particle Theory Group, Department of Physics, College of William and Mary, Williamsburg, VA 23187, USA

(Dated: August 6, 2009)

In the Minimal Supersymmetric Standard Model (MSSM) with three generations of fermions, there is a stringent upper bound on the mass of the lightest neutral Higgs h , and the mass of the charged Higgs H^+ , must be close (within tens of GeV) to the heavier neutral Higgs, H , and the pseudoscalar Higgs A . In this Brief Report, we show that in the four generation MSSM, the upper bound on the h mass is much higher, as high as 400 GeV, and H^+ is generally much heavier than the A , allowing the $H^+ \rightarrow AW^+$ decay, potentially changing search strategies for both the charged Higgs and the pseudoscalar. The H mass, on the other hand, remains within tens of GeV of the charged Higgs mass.

I. INTRODUCTION

One of the most important constraints on supersymmetric models comes from considerations of the mass of the lightest neutral Higgs, h . In the minimal supersymmetric standard model (MSSM), the mass of the h field at tree level [1, 2] must be less than the Z mass, but radiative corrections (primarily due to top quark/squark loops) can lift this to 135 GeV [3]. While non-minimal supersymmetric models can evade this bound somewhat, it is very difficult [4] to raise the upper bound by more than a few tens of GeV.

Another constraint comes from the mass of the charged Higgs boson, H^+ . At tree level, one has [1, 2]

$$M_{H^+}^2 = M_A^2 + M_W^2 \quad (1)$$

where A is the pseudoscalar Higgs mass. Thus, the charged Higgs mass cannot be much heavier than the pseudoscalar, and the decay $H^+ \rightarrow AW^+$ is kinematically forbidden[5]. Similar considerations involving the mass of the heavier neutral scalar, H , show that at tree level

$$M_h^2 + M_H^2 = M_{H^+}^2 + M_Z^2 - M_W^2 \quad (2)$$

which also shows that the charged Higgs cannot decay into HW^+ . Radiative corrections to these formulae have been calculated [6] and, as in the neutral case, are typically tens of GeV.

In this Brief Report, we note that one of the simplest extensions of the MSSM substantially changes these bounds; the addition of a sequential fourth generation. Interest in a fourth chiral generation (with a sufficiently heavy neutrino to avoid contributing to the Z width) has fluctuated over the years [7, 8]. In the early part of this

decade, it was believed by many that electroweak radiative corrections ruled out a chiral fourth generation [9]. However, this belief was based on the assumption that the quark masses were degenerate. It has been shown [10] that relaxing this assumption allows a fourth generation, and for a mass splitting of $O(10)\%$, the S and T corrections fit inside the one-sigma error ellipse [11].

Since corrections to the above mass relations vary as the fourth power of the top quark mass, it is clear that they could be substantially enhanced by a fourth generation. Since the mass splitting between the b' and t' quarks is much smaller than their masses, we will assume that they are degenerate in our analysis. We will consider masses between 250 and 400 GeV. The lower bound is somewhat lower than current Tevatron limits [12], since it might be possible to weaken these limits if supersymmetric decay modes are allowed. The upper bound comes from the requirement that perturbation theory be valid. Generically, the requirement that the Yukawa couplings, g_Y in the MSSM satisfy the condition $g_Y^2/4\pi < 1$ implies [13] that $1/x < \tan\beta < x$, where $x = \sqrt{2\pi(v/M)^2 - 1}$, $\tan\beta$ is the ratio of vacuum expectation values, M is the fermion mass, and $v = 246$ GeV. This window of $\tan\beta$ closes for masses greater than 500 GeV. Since the Yukawa coupling increases with scale, and we want the model to be valid up to at least a TeV, we will set the upper bound to 400 GeV.

In the next section, we look at the lightest neutral scalar, and then consider the mass relations involving the charged Higgs scalar in Section III. Section IV contains our conclusions.

II. THE LIGHTEST NEUTRAL HIGGS MASS

As noted above, $\tan\beta$ must be fairly close to one in order for perturbation theory be valid. In this case, the tree level value of the lightest neutral Higgs mass is negligible, and the entire mass must come from radiative corrections. In the three generation case, these corrections have been calculated, first in Ref. [3], and then

*sean.litsey@yale.edu

†mtsher@wm.edu

with increasing precision by many others [14]. Generally, the contributions from radiative corrections will not be sufficient to increase the mass above 100 GeV (if the tree level mass is negligible).

In the case of four generations, this is not the case. Some early works exploring four generation supersymmetry models[15] did not really address the possibility of fourth generation quarks above 300 GeV. More recently, an exploration of the electroweak phase transition in the four generation MSSM appeared[16], but it used a value of $\tan\beta$ which put the b' Yukawa coupling far outside the region of perturbative validity. Last year, Murdock, et al.[17] discussed ways of extending the validity of perturbation theory in the four generation MSSM. The only recent discussion of the lightest Higgs mass in the four generation MSSM was the work of Fok and Kribs [18], who plotted the lightest Higgs mass as a function of the heavy quark mass. They did restrict the model to the case in which the scalar t' and b' masses were within twenty percent of the t' and b' masses, since that was necessary to obtain sufficient electroweak baryogenesis.

The expression used by Fok and Kribs [18] in the limit in which mixing between scalar quarks is ignored, all squarks have the same mass, the heavier Higgs bosons are all very heavy and $\tan\beta = 1$ is

$$m_h^2 = \sum_{f=t,t',b'} \frac{3}{2\pi^2} \frac{m_f^4}{v^2} \ln \frac{m_{SUSY}^2}{m_f^2} \quad (3)$$

where $v = 246$ GeV. A complete expression which includes two-loop leading logs, as well as squark mixing, can be found in the review article of Carena and Haber [19]. We have used that complete expression, and have plotted the results in Figure 1 for various values of the SUSY scale.

In the graph shown, we have taken the mass of the pseudoscalar to be either 500 or 1000 GeV, and have fixed $\tan\beta = 1$. The results are very insensitive to squark mixing and the precise values of $\tan\beta$ and μ . However, they are sensitive to the pseudoscalar mass. For a pseudoscalar mass of several TeV, the results agree with that given in Eq. (3), but for pseudoscalar masses of 1000 (500) GeV, the results (for $m_{t'} = 400$ GeV) are smaller than that of Eq. (3) by 15 (35) percent.

We see that a large mass for the lightest Higgs boson in the four generation MSSM not only possible, but is likely for SUSY scales above several hundred GeV. Even a mass of close to 400 GeV cannot be excluded.

III. THE CHARGED HIGGS MASS

As was noted earlier, Eq. (1) prevents the decay $H^+ \rightarrow AW^+$ at tree level. This makes detection of the charged Higgs quite difficult, since the $H^+ \rightarrow \bar{t}b, \bar{b}t$ decay has very large backgrounds. Radiative corrections to Eq. (1) can be found in Ref. [6], and when these are modified to account for a fourth generation, the decay may no longer




FIG. 1: Mass of the lightest Higgs in the MSSM as a function of the t' mass. The squark masses are presumed to be equal and with a value of M_{SUSY} . The solid (dashed) lines are for a pseudoscalar mass of 1000 (500) GeV, and the curves, from the lower curve up, correspond to $M_{SUSY} = 600, 800, 1000$ GeV, respectively

be kinematically forbidden. One obtains an expression[6] for $\Delta m \equiv m_{H^+} - m_A$ which depends primarily on $m_{t'}$ and m_{SUSY} . The $\tan\beta$ dependence is very small due to the above requirement of the validity of perturbation theory. The dependence on $m_{t'}$ is illustrated in Figure 2 for several values of m_{SUSY} and $\tan\beta$ fixed to 1. We see that the $H^+ \rightarrow AW^+$ decay becomes kinematically allowed for much of parameter space.

The situation is different with respect to the decay $H^+ \rightarrow HW^+$. Radiative corrections to Eq. (2) are given in Ref. [19], and these can easily be generalized to a fourth generation. We find that m_H is never less than $m_{H^+} - m_W$, and in fact usually $m_H > m_{H^+}$ by tens of GeV. Therefore this decay process is still forbidden, even with higher order corrections.

In addition to the possibility of the decay process $H^+ \rightarrow AW^+$, one has the conventional processes $H^+ \rightarrow \bar{t}b, \bar{b}t$ and, for sufficiently heavy charged Higgs bosons, $H^+ \rightarrow \bar{t}'b', \bar{b}'t'$. Furthermore, one can also have $H^+ \rightarrow hW^+$. The latter decay will be small in this case. The reason is that in the limit of $M_A \gg 100$ GeV, one has[2] $\alpha = -\beta$, and thus the coupling of the light Higgs to the charged Higgs and a W , which is proportional to $\cos(\alpha - \beta)$, vanishes as $\tan\beta \approx 1$. Of course, one might also have various supersymmetric decay modes. We will only consider decays into $\bar{t}b, \bar{t}'b'$ and AW^+ .

In Figure 3, we plot the branching ratios of $H^+ \rightarrow AW^+, H^+ \rightarrow \bar{t}b, \bar{b}t$ and $H^+ \rightarrow \bar{t}'b', \bar{b}'t'$ versus $m_{t'}$, choosing $m_{SUSY} = 1000$ GeV and $m_A = 500$ GeV. Note that




FIG. 2: The mass difference between the charged Higgs scalar and the pseudoscalar in the MSSM as a function of the t' mass. The squark masses are presumed to be equal and with a value of M_{SUSY} . The curves, from the lower curve up, correspond to $M_{SUSY} = 600, 800, 1000$ GeV, respectively. $M(avg)$ is the average of the charged scalar and pseudoscalar masses.




FIG. 3: The branching ratio for $H^+ \rightarrow AW^+$, $H^+ \rightarrow \bar{t}b, \bar{t}t$ and $H^+ \rightarrow \bar{t}'b', \bar{b}'t'$ versus $m_{t'}$, choosing $M_{SUSY} = 1000$ GeV and $M_A = 500$ GeV.

if the pseudoscalar mass is 1000 GeV, then one can see from Figure 2 that $H^+ \rightarrow AW^+$ is only allowed for very large values of $m_{t'}$, and the branching ratio for this mode is always less than a percent. For $m_A = 500$ GeV, the $H^+ \rightarrow \bar{t}'b', \bar{b}'t'$ decay rapidly becomes kinematically inaccessible, and the branching ratio for $H^+ \rightarrow AW^+$ can be as large as 20%. Note that in this case, the pseudoscalar will primarily decay into $\bar{t}t$, and observation of the associated W may be the only way to detect it[21].

IV. CONCLUSIONS

In the MSSM, the Higgs sector is very tightly restricted. There is an upper bound of approximately 130 GeV on the lightest Higgs mass and the charged Higgs mass cannot differ by more than tens of GeV from the pseudoscalar and heavier neutral Higgs masses, thus ruling out two-body decays into these states. In this Brief Report, motivated by recent interest in four generation models, we have considered the effects of radiative correction in the four generation MSSM. The upper bound on the lightest Higgs mass is substantially increased, and can be as large as 400 GeV for reasonable parameter values. The mass splitting between the charged Higgs and pseudoscalar is increased to the point where the $H^+ \rightarrow AW^+$ decay is kinematically allowed, whereas the splitting between the charged Higgs and heavy neutral scalar remains small.

Should a fourth generation exist, it will likely be discovered shortly at the LHC. Our main point is that radiative corrections from these quarks can have major implications for the Higgs sector, not only for the MSSM, but other extensions of the Standard Model.

Acknowledgments

We are grateful to Chris Carone, David Reiss and Graham C. Kribs for useful discussions. This work was supported by the National Science Foundation grant NSF-PHY-0757481 and the REU grant NSF-PHY-0755262.

[1] K. Inoue, A. Kakuto, H. Komatsu and S. Takeshita, Prog. Theor. Phys. **67**, 1889 (1982); R. A. Flores and M. Sher, Annals Phys. **148**, 95 (1983). H. E. Haber and G. L. Kane, Phys. Rept. **117**, 75 (1985).

[2] J. F. Gunion, H. E. Haber, G. L. Kane and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Reading 1990).

[3] H. E. Haber and R. Hempfling, Phys. Rev. Lett **66**, 1815

- (1991); J. R. Ellis, G. Ridolfi and F. Zwirner, Phys. Lett. B **257**, 83 (1991); Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991).
- [4] G. L. Kane, C. F. Kolda and J. D. Wells, Phys. Rev. Lett. **70**, 2686 (1993) [arXiv:hep-ph/9210242].
- [5] For a discussion of the symmetries giving these relations, see J. M. Gerard and M. Herquet, Phys. Rev. Lett. **98**, 251802 (2007) [arXiv:hep-ph/0703051].
- [6] M. A. Diaz and H. E. Haber, Phys. Rev. D **45**, 4246 (1992).
- [7] For a review and an extensive set of references of work before 2000, see P. H. Frampton, P. Q. Hung and M. Sher, Phys. Rept. **330**, 263 (2000) [arXiv:hep-ph/9903387].
- [8] For a review and extensive set of references of work after 2000, see B. Holdom, W. S. Hou, T. Hurth, M. L. Mangano, S. Sultansoy and G. Unel, arXiv:0904.4698 [hep-ph].
- [9] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
- [10] M. Maltoni, V. A. Novikov, L. B. Okun, A. N. Rozanov and M. I. Vysotsky, Phys. Lett. B **476**, 107 (2000) [arXiv:hep-ph/9911535]; H. J. He, N. Polonsky and S. Su, Phys. Rev. D **64**, 053004 (2001) [arXiv:hep-ph/0102144]; B. Holdom, Phys. Rev. D **54**, 721 (1996) [arXiv:hep-ph/9602248];
- [11] G. D. Kribs, T. Plehn, M. Spannowsky and T. M. P. Tait, Phys. Rev. D **76**, 075016 (2007) [arXiv:0706.3718 [hep-ph]]; J. Alwall *et al.*, Eur. Phys. J. C **49**, 791 (2007) [arXiv:hep-ph/0607115].
- [12] www-cdf.fnal.gov/physics/exotic/r2a/20090327.bprime/; www-cdf.fnal.gov/physics/new/top/2008/tprop/Tprime2.8/public.html
- [13] E. De Pree, G. Marshall and M. Sher, arXiv:0906.4500 [hep-ph].
- [14] R. Barbieri and M. Frigeni, Phys. Lett. B **258**, 395 (1991); M. Drees and M. M. Nojiri, Phys. Rev. D **45**, 2482 (1992); J. A. Casas, J. R. Espinosa, M. Quiros and A. Riotto, Nucl. Phys. B **436**, 3 (1995) [Erratum-ibid. B **439**, 466 (1995)] [arXiv:hep-ph/9407389]; J. R. Espinosa and R. J. Zhang, JHEP **0003**, 026 (2000) [arXiv:hep-ph/9912236]; J. R. Espinosa and R. J. Zhang, Nucl. Phys. B **586**, 3 (2000) [arXiv:hep-ph/0003246]; A. Brignole, G. Degrandi, P. Slavich and F. Zwirner, Nucl. Phys. B **631**, 195 (2002) [arXiv:hep-ph/0112177]; D. M. Pierce, J. A. Bagger, K. T. Matchev and R. J. Zhang, Nucl. Phys. B **491**, 3 (1997) [arXiv:hep-ph/9606211]; H. E. Haber, R. Hempfling and A. H. Hoang, Z. Phys. C **75**, 539 (1997) [arXiv:hep-ph/9609331]; S. Heinemeyer, W. Hollik and G. Weiglein, Phys. Lett. B **440**, 296 (1998) [arXiv:hep-ph/9807423]; S. Heinemeyer, W. Hollik and G. Weiglein, Eur. Phys. J. C **9**, 343 (1999) [arXiv:hep-ph/9812472].
- [15] J. F. Gunion, D. W. McKay and H. Pois, Phys. Rev. D **53**, 1616 (1996) [arXiv:hep-ph/9507323]; M. S. Carena, H. E. Haber and C. E. M. Wagner, Nucl. Phys. B **472**, 55 (1996) [arXiv:hep-ph/9512446].
- [16] S. W. Ham, S. K. Oh and D. Son, Phys. Rev. D **71**, 015001 (2005) [arXiv:hep-ph/0411012].
- [17] Z. Murdock, S. Nandi and Z. Tavartkiladze, Phys. Lett. B **668**, 303 (2008) [arXiv:0806.2064 [hep-ph]].
- [18] R. Fok and G. D. Kribs, Phys. Rev. D **78**, 075023 (2008) [arXiv:0803.4207 [hep-ph]].
- [19] M. S. Carena and H. E. Haber, Prog. Part. Nucl. Phys. **50**, 63 (2003) [arXiv:hep-ph/0208209].
- [20] A. G. Akeroyd and S. Baek, Phys. Lett. B **525**, 315 (2002) [arXiv:hep-ph/0105228].
- [21] For a discussion of the phenomenology at the LHC, see S. de Visscher, J. M. Gerard, M. Herquet, V. Lemaitre and F. Maltoni, arXiv:0904.0705 [hep-ph].